

1. Risk factors for loss of residual renal function in children treated with chronic peritoneal dialysis. Ha IS, Yap HK, Munarriz RL, Zambrano PH, Flynn JT, Bilge I, Szczepanska M, Lai WM, Antonio ZL, Gulati A, Hooman N, van Hoeck K, Higuera LM, Verrina E, Klaus G, Fischbach M, Riyami MA, Sahpazova E, Sander A, Warady BA, Schaefer F. *Kidney Int.* 2015. [Epub ahead of print]

2. Interdialytic weight gain in oligoanuric children and adolescents on chronic hemodialysis.

Paglialonga F, Consolo S, Galli MA, Testa S, Edefonti A.

*Pediatr Nephrol.* 2015 Jun;30(6):999-1005.

see associated Editorial: The interdialytic weight gain: a simple marker of left ventricular hypertrophy in children on chronic haemodialysis. By Fischbach M, Zaloszc A, Shroff R.

*Pediatr Nephrol.* 2015 Jun;30(6):859-63.

3. AV fistula creation in paediatric patients: outcome is independent of demographics and fistula type reducing usage of venous catheters. Matoussevitch V, Taylan C, Konner K, Gawenda M, Kuhr K, Hoppe B, Brunkwall J. *J Vasc Access.* 2015.

4. Association of Body Mass Index with Patient-Centered Outcomes in Children with ESRD. Ku E, Glidden DV, Hsu CY, Portale AA, Grimes B, Johansen. *J Am Soc Nephrol.* 2015 [Epub ahead of print]

5. Considerable variations in growth hormone policy and prescription in paediatric end-stage renal disease across European countries-a report from the ESPN/ERA-EDTA registry. van Huis M, Bonthuis M, Sahpazova E, Mencarelli F, Spasojević B, Reusz G, Caldas-Afonso A, Bjerre A, Baiko S, Vondrak K, Molchanova EA, Kolvek G, Zaikova N, Böhm M, Ariceta G, Jager KJ, Schaefer F, van Stralen KJ, Groothoff JW. *Nephrol Dial Transplant.* 2015. [Epub ahead of print]